

AD-A073 918

INDIANA UNIV AT BLOOMINGTON DEPT OF CHEMISTRY  
INTRIGUING ASPECTS OF 1,2-DIALKYL-DIMOLYBDENUM AND -DITUNGSTEN --ETC(U)  
AUG 79 M H CHISHOLM, D A HAITKO

F/G 7/3

N00014-76-C-0826


UNCLASSIFIED

TR-79-4

NL

| OF |


AD  
A073918



END  
DATE  
FILMED

-10-79

DDC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

LEVEL

12  
B.S.

OFFICE OF NAVAL RESEARCH

Contract N00014-76-C-0826

Task No. NR 056-625

TECHNICAL REPORT NO. 79-4

ADA073918

INTRIGUING ASPECTS OF 1,2-DIALKYL-DIMOLYBDENUM AND  
-DITUNGSTEN (M=M) CHEMISTRY,

(M TRIPLE BOND M)

by

Malcolm H. Chisholm and Deborah A. Haitko

Prepared for Publication

in

Journal of the American Chemical Society

DDC  
RECEIVED  
SEP 18 1979  
REGISTERED  
C

Department of Chemistry

Indiana University

Bloomington, Indiana 47405

27 August 27, 1979

12 13p.

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

Approved for Public Release: Distribution Unlimited

79 09 17 072

176 685

DDC FILE COPY

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Intriguing Aspects of 1,2-Dialkyl-dimolybdenum and -ditungsten (M≡M) Chemistry		5. TYPE OF REPORT & PERIOD COVERED Technical Report 1979
		6. PERFORMING ORG. REPORT NUMBER TR-79-4 ✓
7. AUTHOR(s) Malcolm H. Chisholm and Deborah A. Haitko		8. CONTRACT OR GRANT NUMBER(s) N00014-76-C-0826 ✓
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry ✓ Indiana University Bloomington, Indiana 47405		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy		12. REPORT DATE August 27, 1979
		13. NUMBER OF PAGES 14
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release: distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Dimolybdenum Dialkyl, $\beta$ -hydrogen elimination, reductive elimination.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  → The preparation of a series of 1,2-dialkyl compounds of dimolybdenum and ditungsten (M≡M) having the general formula $M_2R_2(NMe_2)_4$ is reported for R = CH <sub>3</sub> , CH <sub>2</sub> CH <sub>3</sub> , CH <sub>2</sub> CD <sub>3</sub> , CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub> , CH(CH <sub>3</sub> ) <sub>2</sub> , CH <sub>2</sub> C(CH <sub>3</sub> ) <sub>3</sub> , CH <sub>2</sub> Si(CH <sub>3</sub> ) <sub>3</sub> and C(CH <sub>3</sub> ) <sub>3</sub> . The isolation of the labelled compounds $M_2(CH_2CD_3)_2(NMe_2)_4$ and the isopropyl and tert-butyl derivatives implies that $\beta$ -hydrogen elimination is either kinetically or thermodynamically not favorable. All of the → beta		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-014-6601



## 20. ABSTRACT continued


above react in hydrocarbon solutions with  $\text{CO}_2$ . When R = a  $\beta$ -hydrogen elimination stabilized alkyl the products are  $\text{M}_2\text{R}_2(\text{O}_2\text{CNMe}_2)_4$  ( $\text{M}=\text{M}$ ) for both  $\text{M} = \text{Mo}$  and  $\text{W}$ . When R contains  $\beta$ -hydrogen atoms, the reaction proceeds differently for molybdenum and tungsten. For molybdenum  $\text{Mo}_2(\text{O}_2\text{CNMe}_2)_4$  ( $\text{M}=\text{M}$ ) is formed along with 1-alkenes and alkanes. Using the labelled compound  $\text{Mo}_2(\text{CH}_2\text{CD}_3)_2(\text{NMe}_2)_4$  this reaction is shown to be intramolecular and to form specifically  $\text{CH}_2=\text{CD}_2$  and  $\text{CH}_2\text{DCD}_3$ , which is consistent with an irreversible  $\beta$ -hydrogen elimination followed by C-H/C-D reductive elimination. Addition of alcohols to  $\text{M}_2\text{R}_2(\text{NMe}_2)_4$  lead to products which depend on the nature of (i) the alcohol, (ii) the alkyl group R and (iii) the metal, Mo or W. For example, the reaction between  $\text{Mo}_2(\text{CH}_2\text{CD}_3)_2(\text{NMe}_2)_4$  and  $\text{Bu}^t\text{OH}$  is shown to give  $\text{Mo}_2(\text{C}_2\text{H}_3\text{D}_2)(\text{OBu}^t)_5$  and  $\text{CH}_2\text{DCD}_3$ .

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or special
A	

## Intriguing Aspects of 1,2-Dialkyl-dimolybdenum and -ditungsten ( $M \equiv M$ ) Chemistry

Sir:

Both historically<sup>1</sup> and commercially<sup>2</sup>  $\sigma$ -alkyl complexes have played a prominent role in the development of mononuclear organotransition metal chemistry. Our syntheses<sup>3</sup> of anti-1,2-dichlorotetradimethylamido compounds,  $M_2Cl_2(NMe_2)_4$  ( $M \equiv M$ ,  $M = Mo, W$ ) affords the opportunity of attempting to prepare 1,2-dialkyl-tetradimethylamido compounds by use of the general metathetic reaction shown in eq. 1. Compounds of the form  $M_2R_2(NMe_2)_4$  allow an investigation of the



reactivity patterns of  $\sigma$ -alkyl groups bonded to the simplest of metal clusters, namely dimetal centers. We wish here to report our extended<sup>4</sup> syntheses of compounds of the general formula  $M_2R_2(NMe_2)_4$  and, in particular, to describe their reactivity patterns towards carbon dioxide and tert-butanol with special attention being given to the labelled compounds  $M_2(CH_2CD_3)_2(NMe_2)_4$ .


From reaction 1 we have obtained  $M_2R_2(NMe_2)_4$  compounds for both  $M = Mo$  and  $W$  where  $R = CH_3, CH_2CH_3, CH_2CD_3, CH_2CH_2CH_2CH_3, CH(CH_3)_2, CH_2CMe_3, CH_2SiMe_3$  and  $CMe_3$ .<sup>5</sup> Purification of these compounds followed the following general procedure: (1) the reaction mixture was warmed slowly to room temperature whereupon the solvent was stripped; (2) hexane was added and the solution was filtered using a medium frit and standard Schlenk-techniques; (3) the hexane solvent was again stripped and the residue was purified by sublimation ( $80-100^\circ C$ ,  $10^{-6}$  cm Hg) to give yellow-orange ( $M=Mo$ ) or orange-red solids ( $M=W$ ). When

R = n-butyl, the compounds were liquids at room temperature and were purified by vacuum distillation.

In all cases the  $^1\text{H}$  nmr spectra of the  $\text{M}_2\text{R}_2(\text{NMe}_2)_4$  compounds obtained in toluene- $d_8$  solution at  $-60^\circ\text{C}$  (100 or 270 MHz) showed a mixture of anti- and -gauche 1,2-dialkyl rotamers. The gauche rotamer was always the predominant rotamer and rather interestingly as the bulk of the alkyl group increases  $\text{CH}_3 \rightarrow \text{CH}_2\text{CMe}_3$  so the predominance of the gauche rotamer increases.<sup>6</sup> At  $-60^\circ\text{C}$  rotations about M-N bonds are restricted leading to proximal and distal N-methyl resonances.<sup>7</sup> A mixture of anti- and -gauche  $\text{M}_2\text{R}_2(\text{NMe}_2)_4$  should have a total of 3 proximal and 3 distal N-methyl resonances.<sup>8</sup>

The thermal stability of these compounds is quite remarkable and the stability of the isopropyl and tert-butyl compounds<sup>9</sup> with respect to isomerization to n-propyl and iso-butyl ligands, respectively, implies that  $\beta$ -hydrogen elimination is either kinetically or thermodynamically not favorable. This is also implied by our observation that when  $\text{LiCH}_2\text{CD}_3$ <sup>10</sup> is used in reaction 1, the resultant 1,2-diethyl compounds retain their  $^2\text{H}$  atoms exclusively in the  $\beta$ -position:  $\text{M}_2(\text{CH}_2\text{CD}_3)_2(\text{NMe}_2)_4$ . Figures 1a and 1b compare the  $^1\text{H}$  nmr spectra of the labelled,  $\text{CH}_2\text{CD}_3$  containing compound and the protio compound, respectively.


In hydrocarbon solutions all of the above compounds react with  $\text{CO}_2$  to give selective insertion into the metal-nitrogen bond.<sup>11</sup> The  $\beta$ -elimination stabilized alkyls of both molybdenum and tungsten react according to eq. 2.



In this reaction the M-M triple bond is retained.<sup>12</sup> However, rather interestingly, the non- $\beta$ -hydrogen elimination stabilized alkyl of dimolybdenum and



ditungsten react quite differently. The molybdenum compounds react according to eq. 3. Reaction of the labelled compound  $\text{Mo}_2(\text{CH}_2\text{CD}_3)_2(\text{NMe}_2)_4$  with  $\text{CO}_2$  in



benzene in a sealed nmr tube led to the specific formation of  $\text{CD}_2=\text{CH}_2$  and  $\text{CH}_2\text{DCD}_3$  as determined by  $^2\text{H}$  nmr. See Figure 2a. The  $^2\text{H}$  spectrum of the ethylene,  $\text{CD}_2=\text{CH}_2$ , showed the predicted splitting based on the reported  $^1\text{H}$ - $^1\text{H}$  couplings reported for ethylene and the known magnetudes of the gyromagnetic ratios of  $^1\text{H}$  and  $^2\text{H}$ .<sup>14</sup> As a further check, we purchased  $^{15}\text{CD}_2=\text{CH}_2$ , trans- $\text{CHD}=\text{CHD}$  and cis- $\text{CHD}=\text{CHD}$  and recorded their  $^2\text{H}$  spectrum in benzene: the observed splittings agreed well with the predicted<sup>14</sup> and thus confirmed unequivocally the formation of exclusively<sup>16</sup>  $\text{CD}_2=\text{CH}_2$  in reaction 3 for  $\text{R} = \text{CH}_2\text{CD}_3$ . In order to check the  $^2\text{H}$  spectrum of  $\text{CH}_2\text{DCD}_3$  in benzene, we prepared a sample by the addition of  $\text{D}_2\text{O}$  to a solid sample of  $\text{LiCH}_2\text{CD}_3$  using a vacuum manifold followed by condensation into an nmr tube. The  $^2\text{H}$  spectrum of the gaseous products showed the same pattern for  $\text{CH}_2\text{DCD}_3$  and, moreover, revealed that some ethylene, specifically  $\text{CD}_2=\text{CH}_2$ , is formed when  $\text{D}_2\text{O}$  is added to anhydrous  $\text{LiCH}_2\text{CD}_3$  in a vacuum manifold.<sup>17</sup>

In a separate experiment, a mixture of  $\text{Mo}_2(\text{CH}_2\text{CH}_3)_2(\text{NMe}_2)_4$  and  $\text{Mo}_2(\text{CH}_2\text{CD}_3)_2(\text{NMe}_2)_4$  in benzene was reacted with  $\text{CO}_2$  in a sealed nmr tube. The  $^2\text{H}$  nmr spectrum was identical to that shown in Figure 2a. Thus within the limits of nmr detection,  $\text{CH}_2\text{DCD}_3$  was the only deuterated ethane present.<sup>16</sup>

When  $\text{CO}_2$  is added to a hydrocarbon solution of  $\text{W}_2(\text{C}_2\text{H}_5)_2(\text{NMe}_2)_4$  a green-blue tungsten containing precipitate is formed and only ethane is evolved.




Using the labelled  $W_2(CH_2CD_3)_2(NMe_2)_4$ , the ethane appears by  $^2H$  nmr spectroscopy to be predominantly  $CH_3CD_3$ . The fate of the ethylene and the nature of tungsten compounds are presently unknown.

Reactions of these  $M_2R_2(NMe_2)_4$  compounds with alcohols are dependent on the nature of (i) the alkyl group R, (ii) the alcohol and (iii) the metal, Mo or W. For example,  $M_2(CH_3)_2(NMe_2)_2$  compounds ( $M = Mo, W$ ) both react with tert-butanol to give  $M_2(CH_3)_2(OBu^t)_4$  compounds. However, addition of  $Pr^iOH$  to  $Mo_2(CH_3)_2(NMe_2)_4$  yields  $Mo_2(OPr^i)_6$  ( $M \equiv M$ ) and  $CH_4$  (2 equiv.). The analogous reaction involving  $W_2(CH_3)_2(NMe_2)_4$  has not yet been carried out but it is known that attempts to prepare  $W_2(OPr^i)_6$  have led to  $W_4(\mu-H)_2(OPr^i)_{14}$ .<sup>18</sup> The reaction of  $Mo_2(C_2H_5)_2(NMe_2)_4$  with  $Bu^tOH$  yielded  $Mo_2Et(OBu^t)_5$  and ethane.<sup>4b</sup> Using the labelled compound  $Mo_2(CH_2CD_3)_2(NMe_2)_4$ , we find that addition of  $Bu^tOH$  causes the exclusive<sup>16</sup> elimination of  $CH_2DCD_3$ . The resulting ethyl ligand in  $Mo_2(C_2H_3D_2)(OBu^t)_5$  contains a statistical distribution of deuteriums on the  $\alpha$  and  $\beta$  carbons, namely integration indicates  $\alpha:\beta = 2:3$ .

The purpose of this communication is to draw attention to the intriguing chemistry associated with alkyl groups coordinated to the dimolybdenum or ditungsten ( $M \equiv M$ ) centers. (1) In  $M_2R_2(NMe_2)_4$  compounds,  $\beta$ -hydrogen elimination is either kinetically or thermodynamically not favorable. (2) Addition of  $CO_2$  to  $Mo_2(CH_2CH_3)_2(NMe_2)_4$  ( $M \equiv M$ ) leads to formation of  $Mo_2(O_2CNMe_2)_4$  ( $M \equiv M$ ) by an intramolecular mechanism. It seems likely that this involves an initial irreversible  $\beta$ -hydrogen elimination which is then followed by C-H reductive elimination. However, it is not known at what step during  $CO_2$  insertion that this process occurs, nor is it known whether C-H reductive elimination occurs across the M-M triple bond,  $H-M \equiv M-Et \rightarrow M \equiv M + Et-H$  or via an oxidative addition-

reductive elimination process at one metal center,  $M^{II} \equiv M^{IV}(H)(Et) \rightarrow M^{II} \equiv M^{II} + Et-H$ . (3) That  $W_2(C_2H_5)_2(NMe_2)_4$  should react so differently with  $CO_2$  is also fascinating, and once again points to the elusive nature of  $W_2(O_2CX)_4$  ( $M \equiv M$ ) compounds.<sup>19</sup> (4) Addition of  $Bu^tOH$  to  $Mo_2(CH_2CD_3)_2(NMe_2)_4$  which leads to  $Mo_2(C_2H_3D_2)(OBu^t)_5$  and  $CH_2DCD_3$  implies that alcoholysis, as with  $CO_2$  insertion (2 above), once again facilitates an irreversible  $\beta$ -hydrogen process which is then followed by C-D reductive elimination. This further implies that formation of the  $\sigma$ -ethyl ligand arises from  $CH_2=CD_2$  and  $Bu^tO-H$ . The statistical distribution of H/D atoms over the  $\alpha$ - and  $\beta$ -carbons in  $\sigma$ -ethyl ligand thus formed is consistent with a then reversible  $\beta$ -hydrogen elimination process favoring the coordinated ethyl ligand:



While all these observations raise intriguing mechanistic questions which cannot be answered at this time, we do note that our observations have at least one parallel with mononuclear transition metal alkyl chemistry: reductive elimination involving C-H bond formation is more facile than reductive elimination involving C-C bond formation.<sup>20</sup>

Malcolm H. Chisholm\*  
Deborah A. Haitko  
Department of Chemistry  
Indiana University  
Bloomington, Indiana 47405

#### Acknowledgements

We thank the Office of Naval Research for support of this work. The  $^1H$  nmr spectrum shown in Figure 1a was obtained through the courtesy of Mr. Dean Katahira at Yale University. Mass spectral data were kindly obtained for us by Mr. Peter Cook and Professor D. C. Bradley at Queen Mary College, London.

### References and Footnotes

1. (a) G. Wilkinson, *Science*, **185**, 109 (1974).  
 (b) M. F. Lappert, P. J. Davidson and R. Pearce, *Acc. Chem. Res.*, **7**, 209 (1974).  
 (c) R. R. Schrock and G. W. Parshall, *Chem. Rev.*, **76**, 243 (1976).
2. J. K. Kochi in "Organometallic Mechanisms and Catalysis", Academic Press, 1978.
3. M. Akiyama, M. H. Chisholm, F. A. Cotton, M. W. Extine and C. A. Murillo, *Inorg. Chem.*, **16**, 2407 (1977).
4. (a) The preparation and structure of  $\text{Mo}_2(\text{CH}_3)_2(\text{NMe}_2)_4$  has been reported: M. H. Chisholm, F. A. Cotton, M. W. Extine and C. A. Murillo, *Inorg. Chem.*, **17**, 2338 (1978).  
 (b) The preparation and reactivity of  $\text{Mo}_2(\text{C}_2\text{H}_5)_2(\text{NMe}_2)_4$  was previously reported: M. H. Chisholm, D. A. Haitko and C. A. Murillo, *J. Am. Chem. Soc.*, **100**, 6262 (1978).
5. Satisfactory analytical data have been obtained. All compounds showed molecular ions in the mass spectrometer,  $\text{M}_2\text{R}_2(\text{NMe}_2)_4^+$ , and we thank Professor D. C. Bradley and Mr. Peter Cook at Queen Mary College for kindly obtaining these valuable spectra for us. This collaboration was promoted by a NATO grant to DCB and MHC.
6. We have no rationale for this observation; our intuition would have suggested the opposite trend.
7. For an assignment of proximal and distal N-Me resonances, see M. H. Chisholm, F. A. Cotton, B. A. Frenz, W. W. Reichert, L. W. Shive and B. R. Stults, *J. Am. Chem. Soc.*, **98**, 4469 (1976).
8. For a detailed discussion, see the variable temperature  $^{13}\text{C}$  nmr studies carried out on  $\text{W}_2(\text{CH}_3)_2(\text{NEt}_2)_4$ : M. H. Chisholm, F. A. Cotton, M. W. Extine, M. Millar and B. R. Stults, *Inorg. Chem.*, **15**, 2244 (1976).
9. Tert-butyl transition metal complexes (and to a lesser extent iso-propyl complexes) are relatively rare. See ref. 2. The reversible  $\beta$ -H-elimination from an alkyl metal complex without attendant reductive elimination of alkane was elegantly studied in alkyl gold(III) chemistry. The isomerization of trans-tert-butyl dimethyl(triphenylphosphine) gold(III) proceeds spontaneously at 25°C in etherial solution to produce the trans-iso-butyl dimethyl(triphenylphosphine) gold(III) complex in quantitative yield. See A. Tamaki and J. K. Kochi, *J.C.S. Chem. Commun.* (1973) 423.



10. Prepared from the reaction between  $\text{CD}_3\text{CD}_2\text{Br}$  and Li shot in hexane; purified by sublimation  $65^\circ\text{C}$ ,  $10^{-6}$  cm Hg.  $\text{CD}_3\text{CH}_2\text{Br}$  was obtained from Merck, Sharp and Dohme.
11. The mechanism of this insertion has been shown to proceed via an amine catalyzed sequence: M. H. Chisholm and M. W. Extine, J. Am. Chem. Soc., 99, 792 (1977).
12. All these compounds show molecular ions in the mass spectrometer and are believed to be directly analogous to the structurally characterized diethylcarbamato analogue  $\text{W}_2(\text{CH}_3)_2(\text{O}_2\text{CNEt}_2)_4$ : M. H. Chisholm, F. A. Cotton, M. W. Extine and B. R. Stults, Inorg. Chem., 16, 603 (1977).
13.  $\text{Mo}_2(\text{O}_2\text{CNMe}_2)_4$  is formed as a fine yellow precipitate in this reaction. It shows a strong molecular ion  $\text{Mo}_2(\text{O}_2\text{CNMe}_2)_4^+$  and a doubly charged molecular ion  $\text{Mo}_2(\text{O}_2\text{CNMe}_2)_4^{2+}$ . In all respects it appears like  $\text{Mo}_2(\text{O}_2\text{CR})_4$  compounds for which scores have been structurally characterized and all of which contain a Mo-Mo quadruple bond.
14. From R. M. Lynden-Bell and N. Sheppard, Proc. Roy. Soc. A., 269, 1385 (1965): for ethylene  $J_{\text{cis}}^{\text{H-H}} = +11.7$  Hz,  $J_{\text{trans}}^{\text{H-H}} = 19.1$  Hz and  $J_{\text{gem}}^{\text{H-H}} = +2.5$  Hz;  $\gamma_{\text{H}} = 2.675 \times 10^4$  radians/sec. gauss,  $\gamma_{\text{H}} = 4.107 \times 10^3$  radians/sec. Using this data it was possible to calculate the splittings of  $\text{CD}_2=\text{CH}_2$ , trans  $\text{CHD}=\text{CHD}$  and cis  $\text{CHD}=\text{CHD}$ . The calculated splittings were obtained by utilizing the program entitled NMRCAL, a part of the Nicolet-80 system. For each of the isomeric dideuteroethylenes a centered three-line pattern was observed with the only apparent distinguishing characteristic being the separation between the outer lines. The calculated versus (observed) splittings in Hertz for each dideuteroethylene are as follows: 1,1-dideuteroethylene 4.8 (4.8); 1,2-cis-dideuteroethylene 3.3 (3.3) and trans 1,2-dideuteroethylene 2.2 (2.2).
15. From Merck, Sharp and Dohme.
16. By exclusively, we mean within the limits of nmr detection which is at least 95%.
17. Alkylolithium and alkylaluminum compounds are well known to undergo  $\beta$ -H-elimination under thermolysis (ref. 2) but we are not aware of anyone reporting the formation of ethylene upon the addition of water. It quite possibly is formed under our conditions because  $\text{LiCH}_2\text{CD}_3$  is present in excess as  $\text{D}_2\text{O}$  is added and therefore thermolysis is a competitive reaction due to local heating effects which accompany hydrolysis. See also B. J. Wakefield in "The Chemistry of Organolithium Compounds", Pergamon Press publisher, 1974, Ch. 15.
18. M. Akiyama, D. Little, M. H. Chisholm, D. A. Haitko, F. A. Cotton and M. W. Extine, J. Am. Chem. Soc., 101, 2504 (1979).







19. A. Bino, F. A. Cotton, Z. Dosi, S. Koch, H. Kuepers, M. Millar and J. C. Sekutowski, *Inorg. Chem.*, 17, 3245 (1978).
20. J. R. Norton, *Acc. Chem. Res.*, 12, 139 (1979) and references therein.


Captions to Figures

Figure 1. Proton NMR spectra recorded at 270 MHz,  $-61^{\circ}\text{C}$  in toluene- $\text{d}_8$  of  
a) \*-anti and gauche -  $\text{Mo}_2(\text{CH}_2\text{CD}_3)_2(\text{NMe}_2)_4$  and b) \*-anti and  
gauche -  $\text{Mo}_2(\text{CH}_2\text{CH}_3)_2(\text{NMe}_2)_4$ . c) \*\* represents residual  
protonated toluene.


Figure 2. Deuterium NMR spectra recorded at  $16^{\circ}\text{C}$  and 33.77 MHz of the gases  
formed in the reactions of a)  $\text{CO}_2$  with  $\text{Mo}_2(\text{CH}_2\text{CD}_3)_2(\text{NMe}_2)_4$  and  
b)  $\text{D}_2\text{O}$  with  $\text{LiCH}_2\text{CD}_3$ . Both spectra were recorded in benzene.  
The scale expansions shown in a) and b) are not the same.



$^2\text{H}$



a)



b)

